

Hitoshi Okamoto M.D., Ph.D**POSITION TITLE:**

Team Leader,
Lab. for Neural Circuit Dynamics of Decision Making
RIKEN Center for Brain Science (CBS)

**ADDRESS:**

2-1 Hirosawa, Wako, Saitama, 351-0198 Japan,
TEL: +81-48-467-9712
FAX: +81-48-467-9714
E-MAIL: hitoshi.okamoto@riken.jp

CV

1983 M.D. Tokyo Univ.
1988 Ph.D. Tokyo Univ. (advisor, Prof. Yoshiki Hotta)
1988~1991 Postdoctoral Research Fellow, Univ. of Michigan, Ann Arbor
(advisor, Prof. John Y. Kuwada)
1991~1993 Assistant Professor, National Institute for Basic Biology (NIBB)
1993~1997 Lecturer and Associate Professor, Dept. of Physiology,
Keio Univ., School of Med. and Adjunct Associate Professor, NIBB
1997~2018, Laboratory Head, Lab. for Developmental Gene Regulation,
RIKEN Brain Science Institute (BSI)
2004~2008, Group Director, Neural Growth and Regeneration Research Group, BSI
2008~2018, Deputy Director of BSI,
2009~2018, Senior Team Leader, BSI
2018~present, Team Leader,
Lab. for Neural Circuit Dynamics of Decision Making
Center for Brain Science (CBS)
Director, RIKEN-Kao Collaboration Center
Adjunct Professor of Waseda Univ., Keio Univ., Tokyo Univ.

Prize

Tokizane Prize (2014)

Publications

- 1) Okamoto, H., Hiromi, Y., Ishikawa, E., Yamada, T., and Hotta, Y. (1985). Molecular analyses of *Drosophila melanogaster* mutant actin genes. pp75-84. in Proceedings of the 11th International Symposium, Division of Biophysics, the Taniguchi Foundation.
- 2) Hiromi, Y., Okamoto, H., Gehring, W. J., and Hotta, Y. (1986). Germline transformation with *Drosophila melanogaster* mutant actin genes induces constitutive expression of heat shock genes. **Cell**, 44:293-301.
- 3) Okamoto, H., Hiromi, Y., Ishikawa, E., Yamada, T., Isoda, K., Maekawa, H., and Hotta, Y. (1986). Molecular characterization of mutant actin genes which induce heat-shock proteins in *Drosophila* flight muscles. **EMBO J.**, 5:589-596.
- 4) Okamoto, H., Sakai, K., Goto, S., Takasu-Ishikawa, E., and Hotta, Y. (1987). Isolation of *Drosophila* genomic clones homologous to the eel sodium channel gene. **Proceedings of the Japan Academy**. 63, SerB:284-288.
- 5) Yoshihara, M. Takasu-Ishikawa, E., Sakai, Y., Hotta, Y., and Okamoto, H. (1988). Single P-element insertional mutagenesis in *Drosophila melenogaster*. **Proceedings of the Japan Academy**. 64, SerB: 172-176.
- 6) Sakai, K., Okamoto, H., and Hotta, Y. (1989). Pharmacological characterization of sodium channels in the primary culture of individual *Drosophila* embryos: neurons of a mutant deficient in a putative sodium channel gene. **Cell Diff. Devel. (Mech. Devel.)** 26:107-118.
- 7) Kuwada, J. Y., Okamoto, H., Bernhardt, R., and Chitnis, A. J. (1990). Growth cone guidance in a simple vertebrate nervous system. pp5-16. in Advance in Neural Regeneration Research. Wiley-Liss, Inc.
- 8) Sakai, Y., Okamoto, H., Mogami, K. Yamada, T., and Hotta, Y. (1990). Actin with tumor related mutation is antimorphic in *Drosophila* muscle: Two distinct modes of myofibrillar disruption by antimorphic actins. **J. Biochem.**, 107: 499-505.
- 9) Okamoto, H., and Kuwada, J. Y. (1991). Outgrowth by fin motor axons in wild type and a finless mutant of the Japanese Medaka fish. **Dev. Biol.**, 146, 49-61.
- 10) Okamoto, H., and Kuwada, J. Y. (1991). Alteration of pectoral fin nerves following ablation of fin buds and by ectopic fin buds in the Japanese Medaka fish. **Dev. Biol.**,

146:62-71.

- 11) Sakai, Y., Okamoto, H., Mogami, K., Matsuo, H., and Hotta, Y. (1991). Heat shock gene activation by mutant actin is independent of myofibril degeneration in Drosophila muscle. **J. Biochem.**, 109: 670-673.
- 12) Inoue, A., Takahashi, M., Hatta, K., Hotta, Y., and Okamoto, H. (1994). Developmental regulation of Isl-1 mRNA expression during neuronal differentiation in embryonic zebrafish. **Dev. Dynam.**, 199:1-11.
- 13) Kuo, C.-H., Uetsuki, T., Kim, C.-H., Tanaka, H., Li, B.-S., Taira, E., Higuchi, H., Okamoto, H., Yoshikawa, K., and Miki, N., (1995). Determination of a necidin cis-acting element required for neuron specific expression by using zebrafish. **Biochem. Biophys. Res. Comm.**, 211:438-446.
- 14) Tokumoto, M., Gong,Z., Tsubokawa, T., Hew,C., Hotta, Y. Uyemura,K., and Okamoto, H. (1995). Molecular heterogeneity among primary moto-neurons and within myotomes revealed by differential mRNA expression of novel Isl-1 homologs in embryonic zebrafish. **Dev. Biol.**, 171:578-589.
- 15) Okamoto, H., Takahashi, M., Inoue, A., Gong, Z., Tsubokawa, T., Hew, C., Uyemura, K., and Hotta, Y. (1996). Isl-1 family in motoneuron specification and target recognition in embryonic zebrafish. pp401-412 in Basic Neuroscience in Invertebrate, Business Center for Academic Societies Japan, Ed. H. Koike.
- 16) Shimoda, N., Chevrette, M., Ekker, M., Kikuchi, Y., Hotta, Y., and Okamoto, H. (1996). Mermaid: a family of short interspersed repetitive elements widespread in vertebrates. **Biochem. Biophys. Res. Comm.**, 220:226-232.
- 17) Shimoda, N., Chevrette, M., Ekker, M., Kikuchi, Y., Hotta, Y., and Okamoto, H. (1996). Mermaid, a family of short interspersed repetitive element, is useful for the zebrafish genome mapping. **Biochem. Biophys. Res. Comm.**, 220:233-237.
- 18) Kim, C-H., Taira, E., Kuo, C-H., Li, B-S., Okamoto, H., Nakahira, K., Ikenaka, K., Higuchi, H. and Miki, N. (1996). Neuron-specific expression of a chicken gicerin cDNA in transient transgenic zebrafish. **Neurochem. Res.**, 22:231-137.
- 19) Kikuchi, Y., Segawa, H., Tokumoto, M., Tsubokawa, T., Hotta, Y., Uyemura, K., and Okamoto, H.(1997). Ocular and cerebellar defects in zebrafish induced by

overexpression of the LIM domain of the Islet-3 LIM/homeodomain protein. **Neuron**, 18:369-382.

- 20) Kikuchi, Y., Segawa, H., Tokumoto, M., Tsubokawa, T., Hotta, Y., Uyemura, K., and Okamoto, H. (1997). Functional analysis of Islet-3 LIM/homeodomain protein in zebrafish brain development. pp73-87, in Molecular Basis of Axon Growth and Nerve Pattern Formation (Taniguchi Symposium on Brain Science No. 20) Fijisawa, H. ed. Japan Scientific Societies Press, Tokyo.
- 21) Higashijima, S., Nose, A., Eguchi, G., Hotta, Y., and Okamoto, H. (1997). Mindin/F-spondin family: novel ECM proteins expressed in the zebrafish embryonic axis. **Dev. Biol.**, 192:211-227.
- 22) Higashijima, S., Okamoto, H., Ueno, N., Hotta, Y., and Eguchi, G. (1997). High-frequency generation of transgenic zebrafish which reliably express GFP in whole muscles or the whole body by using promoter of zebrafish origin. **Dev. Biol.**, 192:289-299.
- 23) Kikuchi, Y., Segawa, H., Tokumoto, Y., Hotta, Y., Uyemura, K., and Okamoto, H. (1999). Islet-3 in the regional specification of the brain. pp108-114, in Neural Development, Uyemura, K., Kawakami, K. and Yazaki eds., Springer.
- 24) Izsvak, Z., Ivics, Z., Shimoda. N., Mohn, D., Okamoto, H., Hackett, P. B. (1999). Short inverted-repeat transposable elements in teleost fish and implications for a mechanism of their amplification. **J. Mol. Evol.**, 48:13-21.
- 25) Mieda, M., Kikuchi, Y., Hirate, Y., Aoki, M., and Okamoto, H. (1999). Compartmentalized expression of zebrafish *ten-m3* and *ten-m4*, homologues of the *Drosophila ten^m/odd Oz* gene, in the central nervous system. **Mech. Dev.**, 67: 223-227.
- 26) Feinstein, Y., Borrell, V., Garcia, C., Burstyn-Cohen, T., Tzarfaty, V., Frumkin, A., Nose, A., Okamoto, H., Higashijima, S., Soriano, E., and Klar, A (1999). F-spondin and mindin: two structurally and functionally related genes expressed in the hippocampus that promote outgrowth of embryonic hippocampal neurons. **Development**, 126: 3637-3648.
- 27) Higashijima, S., Hotta, Y., and Okamoto, H. (2000). Visualization of cranial motor neurons in live transgenic zebrafish expressing GFP under the control of the Islet-1 promoter/enhancer. **J. Neurosci.**, 20:206-218.

- 28) Saito, T., Hascilowicz, T., Ohkido, I., Kikuchi, Y., Okamoto, H., Hayashi, S., Murakami, Y., and Matsufuji, S. (2000). Two zebrafish (*Danio rerio*) antizymes with expression and activities. **Biochem. J.**, 345: 99-106.
- 29) Masai, I., Okamoto, H., and Wilson, S. W. (2000). Midline signals regulate retinal neurogenesis in zebrafish. **Neuron**, 27:251-263.
- 30) Yeo, S.-Y., Little, M. H., Yamada, T., Miyashita, T., Halloran M. C, Kuwada, J. Y., Huh T.-L., Okamoto, H. (2001). Overexpression of a Slit Homologue Impairs Convergent Extension of the Mesoderm and Causes Cyclopia in Embryonic Zebrafish. **Dev. Biol.**, 230:1-17.
- 31) Segawa, H., Miyashita, T., Hirate, Y., Higashijima, S., Chino, N., Uyemura, K., Kikuchi, Y. and Okamoto, H. (2001). Functional repression of Islet-2 by disruption of the heteromeric complex with Ldb impairs peripheral axonal outgrowth by the primary sensory and motor neurons in embryonic zebrafish. **Neuron**, 30:423-436.
- 32) Mizuno, T., Kawasaki, M., Nakahira, M., Kagamiyama, H., Kikuchi, Y., Okamoto, H., Mori, K., and Yoshihara, Y. (2001). Molecular diversity in zebrafish ncam family: three members with different vase usage and distinct localization. **Mol. Cell. Neurosci.**, 18:119-130.
- 33) Ando, H., Furuta, T., Tsien, R. Y. and Okamoto, H. (2001). Photo-mediated gene activation using caged RNA/DNA in zebrafish embryos. **Nature Genetics**, 28:317-325.
- 34) Hirate, Y., Mieda, M., Harada, T., Yamasu, K. and Okamoto, H. (2001). Identification of *ephrin-A3* and novel genes specific to the midbrain-MHB in embryonic zebrafish by ordered differential display. **Mech. Dev.**, 107: 83-96.
- 35) Zeller, J., Schneider, V., Malayaman, S., Higashijima, S., Okamoto, H., Gui, J., Lin, S., and Granato, M. (2002). Migration of zebrafish spinal motor nerves into the periphery requires multiple myotome-derived cues. **Dev. Biol.**, 252:241-56
- 36) Bingham, S., Higashijima, S., Okamoto, H., and Chandrasekhar, A. (2002). The Zebrafish trilobite gene is essential for tangential migration of branchiomotor neurons. **Dev Biol.**, 242:149-60.
- 37) Okamoto., H., Segawa, H., and Higashijima, S. (2003). Toward genetic dissection of motor neuron differentiation. Aquatic Genome-Steps toward a great future., Shimizu,

N., Aoki, T., Takashima, F., eds., Springer-Verlag, Tokyo.

- 38) Hirate, Y., Okamoto, H., and Yamasu, K. (2003). Structure of the zebrafish fasciclin I-related extracellular matrix protein (big-h3) and its characteristic expression during embryogenesis. **Gene Expr Patterns.**, 3:331-6.
- 39) Ando, H., and Okamoto, H. (2003). Practical procedures for ectopic induction of gene expression in zebrafish embryos using Bhc-diazo-caged mRNA. **Methods Cell Sci.**, 25:25-31.
- 40) Hutson, L.D., Juryne, M.J., Yeo, S.Y., Okamoto, H., and Chien, C.B. (2003). Two divergent slit1 genes in zebrafish. **Dev. Dyn.**, 228:358-369.
- 41) Masai, I., Lele, Z., Yamaguchi, M., Komori, A., Nakata, A., Nishiwaki, Y., Wada, H., Tanaka, H., Nojima, Y., Hammerschmidt, M., Wilson, S.W., and Okamoto, H. (2003). N-cadherin mediates retinal lamination, maintenance of forebrain compartments and patterning of retinal neurites. **Development**, 130:2479-2494.
- 42) Okamoto, H., Hirate, Y., and Ando, H. (2004). Systematic identification of factors in zebrafish regulating the early midbrain and cerebellar development by ordered differential display and caged mRNA technology. **Front. Biosci.**, 9:93-99.
- 43) Miyashita, T., Yeo, S.Y., Hirate, Y., Segawa, H., Wada, H., Little, M.H., Yamada, T., Takahashi, N., and Okamoto, H. (2004). PlexinA4 is necessary as a downstream target of Islet2 to mediate Slit signaling for promotion of sensory axon branching. **Development**, 131:3705-3715.
- 44) Yeo, S.Y., Miyashita, T., Fricke, C., Little, M.H., Yamada, T., Kuwada, J.Y., Huh, T.L., Chien, C.B., and Okamoto, H. (2004). Involvement of Islet-2 in the Slit signaling for axonal branching and defasciculation of the sensory neurons in embryonic zebrafish. **Mech. Dev.**, 121:315-324.
- 45) Hanaoka, R., Ohmori, Y., Uyemura, K., Hosoya, T., Hotta, Y., Shirao, T., and Okamoto H. (2004). Zebrafish *gcmb* is required for pharyngeal cartilage formation. **Mech. Dev.**, 121:1235-1247
- 46) Ando H, Furuta T, and Okamoto H. (2004) Photo-mediated gene activation by using caged mRNA in zebrafish embryos. **Methods Cell Biol.**, 77:159-71.

- 47) Emoto, Y., Wada, H., Okamoto, H., Kudo A., and Imai, Y. (2005). Retinoic acid-metabolizing enzyme Cyp26a1 is essential for determining territories of hindbrain and spinal cord in zebrafish, **Dev. Biol.**, 278:415-427.
- 48) Uemura, O., Okada, Y., Ando, H., Guedj, M., Higashijima, S., Shimazaki, T., Chino, N., Okano, H., and Okamoto, H. (2005). Comparative functional genomics revealed conservation and diversification of three enhancers of the *isl1* gene for motor and sensory neuron-specific expression, **Dev. Biol.**, 278:567-606.
- 49) Aizawa, H., Bianco, I.H., Hamaoka, T., Miyashita, T., Uemura, O., Concha, M.L., Russell, C., Wilson, S.W., and Okamoto, H., (2005). Laterotopic Representation of Left-Right Information onto the Dorso-Ventral Axis of a Zebrafish Midbrain Target Nucleus, **Curr. Biol.**, 15: 238-243,
- 50) Miyasaka, N., Sato, Y., Yeo, S-Y., Chien, C-B., Okamoto, H., and Yoshihara, Y. (2005). Robo2 mediates the formation of an initial axon scaffold essential for establishment of precise glomerular map in the zebrafish olfactory system. **Development**, 132:1283-93.
- 51) Kawakami, A., Nojima, Y., Toyoda, A., Takahoko, M., Satoh, M., Tanaka, H., Wada, H., Masai., I., Terasaki, H., Sakaki, Y., Takeda, H., and Okamoto, H. (2005). The zebrafish-secreted matrix protein You/Scube2 is implicated in long-range regulation of Hedgehog signaling. **Curr Biol.**, 15: 480-488
- 52) Masai, I., Yamaguchi, M., Tonou-Fujimori, N., Komori, A., and Okamoto, H. (2005). Requirement of Hedgehog signalling for the cell-cycle exit of retinal progenitor cells in zebrafish. **Development**, 132:1539-1553
- 53) Li, Q., Shirabe, H., Thisse, C., Thisse, B., Okamoto, H., Masai,I., and Kuwada, J.Y. (2005). Chemokine signaling guides axons within the retina in zebrafish, **J. Neurosci.**, 25:1711-1717.
- 54) Wada, H., Iwasaki, M., Sato, T., Masai, I., Nishiwaki, Y., Tanaka, H., Sato, A., Nojima, Y., and Okamoto, H. (2005). Dual roles of zygotic and maternal Scribble1 in neural migration and convergent extension movements in zebrafish embryos. **Development**, 132:2273-2285.
- 55) Yamaguchi, M., Tonou-Fujimori, N., Komori, A., Maeda, R., Nojima, Y., Li, H., and Okamoto., H., and Masai, I. (2005). Histone deacetylase 1 regulates retinal

neurogenesis in zebrafish by suppressing Wnt and Notch signaling pathways, **Development**, 132:3027-43

- 56) Ando, H., Kobayashi, M., Tsubokawa, T., Uyemura, K., Furuta, T., and Okamoto, H. (2005). Lhx2 mediates the activity of Six3 in zebrafish forebrain growth. **Dev. Biol.**, 287:456-468.
- 57) Hirate, Y., and Okamoto, H. (2006) Canopy1, a Novel Regulator of FGF Signaling around the Midbrain-Hindbrain Boundary in Zebrafish. **Curr. Biol.**, 16:421-7.
- 58) Sato, T., Takahoko, M., and Okamoto, H. (2006). HuC:Kaede, a useful tool to label neural morphologies in networks in vivo. **Genesis**, 44:136-42.
- 59) Wada, H., Tanaka, H., Nakayama, S., Iwasaki, M., and Okamoto, H. (2006). Frizzled3a and Celsr2 regulate neuroepithelial exclusion of migrating facial motor neurons in the zebrafish hindbrain. **Development**, 133:4749-59.
- 60) Aoki, M., Mieda, M., Ikeda, T., Hamada, Y., Nakamura, H., and Okamoto, H. (2007). R-spondin3 is required for mouse placental development. **Dev. Biol.**, 301:218-226.
- 61) Aizawa, H., Goto, M., and Okamoto, H. (2007). Temporally regulated asymmetric neurogenesis causes left-right difference in the zebrafish habenular structures. **Dev. Cell**, 12:87-98.
- 62) Sassa, T., Aizawa, H., and Okamoto H., (2007) Visualization of two distinct classes of dorsal commissural neurons by *gad2* and *zic1* promoter/enhancer elements in developing zebrafish revealed the neuronal connectivity related to the auditory and lateral line system. **Dev. Dyn.**, 236:706-718.
- 63) Sato, T., Hamaoka, T., Aizawa, H., Hosoya, T., and Okamoto, H. (2007). Genetic single-cell mosaic analysis implicates ephrinB2 reverse signaling in projections from the posterior tectum to the hindbrain in zebrafish. **J. Neurosci.**, 27:5271-9.
- 64) Tanaka, H., Maeda, R., Shoji, W., Wada, H., Masai, I., Shiraki, T., Kobayashi, M., Nakayama, R., and Okamoto H. (2007). Novel mutations affecting axon guidance in zebrafish and a role for plexin signaling in the guidance of trigeminal and facial nerve axons. **Development**, 134:3259-3269.

- 65) Okamoto H. (2007) Yin-yang ways of controlling gene expression are now in our hands. **ACS Chem. Biol.**, 2:646-648.
- 66) Omata, Y., Nojima, Y., Nakayama, S., Okamoto, H., Nakamura, H., and Funahashi, J. (2007) Role of Bone morphogenetic protein 4 in zebrafish semicircular canal development. **Dev. Growth. Differ.**, 49:711-719.
- 66) Aoki, M., Kiyonari, H., Nakamura H., and Okamoto H. (2008). R-spondin2 expression in the apical ectodermal ridge is essential for outgrowth and patterning in mouse limb development. **Dev. Growth. Differ.**, 50:85–95.
- 67) Yamaguchi, M., Fujimoto-Tonou, N., Yohimura, Y., Kishi, T., Okamoto, H., and Masai, I. (2008). Mutation of DNA primase causes extensive apoptosis of retinal neurons through the activation of DNA damage checkpoint and tumor suppressor p53, **Development**, 135:1247-57.
- 68) Okamoto, H., Sato, T., and Aizawa H., (2008). Transgenic technology for visualization and manipulation of the neural circuits controlling behavior in zebrafish, **Dev. Growth. Differ.**, 50 Suppl 1:S167-75.
- 69) Nishiwaki, Y., Komori, A., Sagara, H., Suzuki, E., Manabe, T., Hosoya, T., Nojima, Y., Wada, H., Tanaka, H., Okamoto, H., and Masai I. (2008). Mutation of cGMP phosphodiesterase 6alpha'-subunit gene causes progressive degeneration of cone photoreceptors in zebrafish. **Mec. Dev.**, 125:932-946.
- 70) Okamoto, H. (2008). An interview with Hitoshi Okamoto, M.D., Ph.D. Interview by Stephen C. Ekker. **Zebrafish**, 5:155-158.
- 71) Carreira-Barbosa, F., Kajita, M., Morel, V., Wada, H., Okamoto, H., Martinez-Arias, A., Fujita, Y., Wilson, S.W., and Tada, M. (2009). Flamingo regulates epiboly and convergence/extension movements through cell cohesive and signalling functions during zebrafish gastrulation. **Development**, 136:383-392.
- 72) Miyasaka, N., Morimoto, K., Tsubokawa, T., Higashijima, S., Okamoto, H., and Yoshihara Y. (2009). From the olfactory bulb to higher brain centers: genetic visualization of secondary olfactory pathways in zebrafish. **J. Neurosci.**, 29:4756-4767.
- 72) Ohata, S., Kinoshita, S., Aoki, R., Tanaka, H., Wada, H., Tsuruoka-Kinoshita, S., Tsuboi, T., Watabe, S., and Okamoto, H. (2009). Neuroepithelial cells require

fucosylated glycans to guide the migration of vagus motor neuron progenitors in the developing zebrafish hindbrain **Development**, 136:1653-1663.

73) Wada, H., and Okamoto, H. (2009). Roles of planar cell polarity pathway genes for neural migration and differentiation. **Dev. Growth. Differ.**, 51:233-240.

74) Wada, H., Okamoto, H. (2009) . Roles of noncanonical Wnt/PCP pathway genes in neuronal migration and neurulation in zebrafish. **Zebrafish**, 6:3-8.

75) Sugiyama, M., Sakaue-Sawano, A., Iimura, T., Fukami, K., Kitaguchi, T., Kawakami, K., Okamoto, H., Higashijima, S., and Miyawaki, A. (2009). Illuminating cell-cycle progression in the developing zebrafish embryo. **Proc. Natl. Acad. Sci. USA.**, 106:20812-7.

76) Amo, R., Aizawa, H., Takahoko, M., Kobayashi, M., Takahoko, R., Aoki, T., and Okamoto, H. (2010). Identification of the zebrafish ventral habenula as a homologue of the mammalian lateral habenula. **J. Neurosci.**, 30:1566-1574.

77) Ito, Y., Tanaka, H., Okamoto, H., and Ohshima, T. (2010). Characterization of neural stem cells and their progeny in the adult zebrafish optic tectum. **Dev. Biol.**, 342:26-38.

78) Okamoto, H., and Ishioka, A. (2010). Zebrafish research in Japan and the National BioResource Project. **Exp. Anim.**, 59:9-12.

79) Palmesino, E., Rousso, D.L., Kao, T.J., Klar, A., Laufer, E., Uemura, O., Okamoto, H., Novitch, B.G., and Kania, A. (2010). Foxp1 and lhx1 coordinate motor neuron migration with axon trajectory choice by gating Reelin signalling. **PLoS Biol.**, 8(8):e1000446.

80) Agetsuma, M., Aizawa, H., Aoki, T., Nakayama, R., Takahoko, M., Goto, M., Sassa, T., Amo, R., Shiraki, T., Kawakami, K., Hosoya, T., Higashijima, S., and Okamoto, H. (2010). The habenula is crucial for experience-dependent modification of fear responses in zebrafish. **Nat. Neurosci.**, 13:1354-1356.

81) Tanaka, H., Nojima, Y., Shoji, W., Sato, M, Nakayama, R, Ohshima, T., and Okamoto, H. (2011). Islet1 selectively promotes peripheral axon outgrowth in Rohon-Beard primary sensory neurons. **Dev. Dyn.**, 240:9-22.

- 82) Ohata, S., Aoki, R., Kinoshita, S., Yamaguchi, M., Tsuruoka-Kinoshita, S., Tanaka, H., Wada, H., Watabe, S., Tsuboi, T., Masai, I. and Okamoto H. (2011). Dual Roles of Notch in Regulation of Apically Restricted Mitosis and Apicobasal Polarity of Neuroepithelial Cells. **Neuron**, 69:215-230.
- 83) Okamoto, H., Agetsuma, M, and Aizawa H. (2012). Genetic dissection of the zebrafish habenula, a possible switching board for selection of behavioral strategy to cope with fear and anxiety. **Dev. Neurobiol.** 72:386-394.
- 84) Matsui, T., Thitamadee, S., Murata, T., Kakinuma, H., Nabetani, T., Hirabayashi, Y., Hirate, Y., and Okamoto H, Bessho Y. (2011). Canopy1, a positive feedback regulator of FGF signaling, controls progenitor cell clustering during Kupffer's vesicle organogenesis. **Proc. Natl. Acad. Sci. USA.**, 108:9881-9886.
- 85) Aizawa, H., Amo, R., and Okamoto, H. (2011). Phylogeny and ontogeny of the habenular structure. **Front Neurosci.** 5:138. doi: 10.3389/fnins.2011.00138.
- 86) Okamoto, H., Agetsuma, M., and Aizawa, H. (2012). Genetic dissection of the zebrafish habenula, a possible switching board for selection of behavioral strategy to cope with fear and anxiety. **Dev. Neurobiol.**, 72:386-394.
- 87) Aizawa, H., Kobayashi, M., Tanaka, S., Fukai, T., and Okamoto, H. (2012). Molecular characterization of the subnuclei in rat habenula. **J. Comp. Neurol.**, 520:4051-4066.
- 88) Aizawa, H., Yanagihara, S., Kobayashi, M., Niisato, K., Takekawa, T., Harukuni, R., McHugh, T.J., Fukai, T., Isomura, Y., and Okamoto, H. (2013). The synchronous activity of lateral habenular neurons is essential for regulating hippocampal theta oscillation. **J. Neurosci.**, 33:8909-8921.
- 89) Okamoto, H., and Aizawa, H. (2013). Fear and anxiety regulation by conserved affective circuits. **Neuron**, 78:411-413.
- 90) Aoki, T., Kinoshita, M., Aoki, R., Agetsuma, M., Aizawa, H., Yamazaki, M., Takahoko, M., Amo, R., Arata, A., Higashijima, S, Tsuboi, T., and Okamoto H. (2013). Imaging of Neural Ensemble for the Retrieval of a Learned Behavioral Program. **Neuron**, 78:881-94.
- 91) Mizuno, H., Sassa, T., Higashijima, S., Okamoto, H., and Miyawaki A. (2013). Transgenic zebrafish for ratiometric imaging of cytosolic and mitochondrial Ca²⁺ response in teleost embryo. **Cell Calcium**, 54:236-245.

- 92) Campbell D.S., and Okamoto, H. (2013). Local caspase activation interacts with Slit-Robo signaling to restrict axonal arborization. **J. Cell Biol.**, 203:657-672.
- 93) Aoki, M., Segawa, H., Naito, M., and Okamoto, H. (2014). Identification of possible downstream genes required for the extension of peripheral axons in primary sensory neurons. **Biochem. Biophys. Res. Commun.**, 445:357-362.
- 94) Nishiya, N., Oku, Y., Kumagai, Y., Sato, Y., Yamaguchi, E., Sasaki, A., Shoji, M., Ohnishi, Y., Okamoto, H., and Uehara, Y. (2014). A zebrafish chemical suppressor screening identifies small molecule inhibitors of the Wnt/β-catenin pathway. **Chem. Biol.**, 21:530-540.
- 95) Okamoto, H. (2014). Neurobiology: sensory lateralization in the fish brain. **Curr. Biol.**, 24:R285-287.
- 96) Okamoto, H. (2014). Minicerebellum, now available for reductionists' functional study. **Proc. Natl. Acad. Sci. USA.**, 111:11580-11581.
- 97) Amo, R., Fredes, F., Kinoshita, M., Aoki, R., Aizawa, H., Agetsuma, M., Aoki, T., Shiraki, T., Kakinuma, H., Matsuda, M., Yamazaki, M., Takahoko, M., Tsuboi, T., Higashijima, S., Miyasaka, N., Koide, T., Yabuki, Y., Yoshihara, Y., Fukai, T. and Okamoto, H. (2014). The habenulo-raphe serotonergic circuit encodes an aversive expectation value essential for adaptive active avoidance of danger. **Neuron**, 84:1034-1048.
- 98) Elbaz, I., Lerer-Goldshtein, T., Okamoto, H., and Appelbaum, L. (2015). Reduced synaptic density and deficient locomotor response in neuronal activity-regulated pentraxin 2a mutant zebrafish. **FASEB J.**, 29:1220-1234.
- 99) Mieda, M., Ono, D., Hasegawa, E., Okamoto, H., Honma, K., Honma, S., Sakurai, T. (2015). Cellular clocks in AVP neurons of the SCN are critical for interneuronal coupling regulating circadian behavior rhythm. **Neuron**, 85:1103-1116.
- 100) Hisano, Y., Sakuma, T., Nakade, S., Ohga, R., Ota, S., Okamoto, H., Yamamoto, T., and Kawahara, A. (2015). Precise in-frame integration of exogenous DNA mediated by CRISPR/Cas9 system in zebrafish. **Sci. Rep.**, 5:8841. doi: 10.1038/srep08841.

- 101) Aoki, R., Tsuboi, T., and Okamoto, H. (2015). Y-maze avoidance: an automated and rapid associative learning paradigm in zebrafish. **Neurosci. Res.**, 91:69-72.
- 102) Chou, M., Amo, R., Kinoshita, M., Cherng, B., Shimazaki, H., Agetsuma, M., Shiraki, T., Aoki, T., Takahoko, M., Yamazaki, M., Higashijima, S., and Okamoto, H. (2016). Social conflict resolution regulated by two dorsal habenular subregions in zebrafish. **Science**, 352:87-90
- 103) Okamoto, H. (2016). For Motor Adjustments, Serotonin Steps In. **Cell**. 167:886-887.
- 104) Mieda, M., Okamoto, H., and Sakurai, T. (2016). Manipulating the Cellular Circadian Period of Arginine Vasopressin Neurons Alters the Behavioral Circadian Period. **Curr. Biol.**, 26:2535-2542.